

PRESCRIPTION TREATMENT[®] brand

TRI-DIE[®]

Silica & Pyrethrum Dust

KILLS: Ants, Bedbugs, Bees, Booklice, Boxelder Bugs, Carpet Beetles, Carpenter Ants, Centipedes, Clover Mites, Cluster Flies, Crickets, Cockroaches, Drywood Termites, Earwigs, Elmleaf Beetles, Firebrats, Fleas, Ground Beetles, Lice, Millipedes, Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders, Stored Product Pests, Powder Post Beetles, Ticks and Wasps,

FOR USE IN AND AROUND: Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals, Hotels, Food Processing Plants, Motels, Nursing Homes, Office Buildings, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes*), Utilities, Warehouses, and other Commercial and Industrial Buildings. *Do not use in aircraft cabins.

Not for use in USDA Meat and Poultry Plants.
For Use by Commercial Applicators.

ACTIVE INGREDIENTS:

Pyrethrins	1.0%
Piperonyl Butoxide, Technical*	10.0%
Amorphous Silica	40.0%

INERT INGREDIENTS:**

* Equivalent to 8.0% (butylcarbityl) (6-propylpiperonyl) ether and 2.0% related compounds.

** Contains Petroleum Distillate

EPA Reg. No. 499-429 EPA Est. No. 499-MO-1^A
EPA Est. No. 67517-MO-1^B
(See code in lot number.)

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

INDOOR TREATMENTS

CRAWLING INSECTS: Ants, Booklice, Carpet Beetles, Crickets, Cockroaches, Scorpions, Silverfish, Firebrats and Spiders: Apply a small amount of TRI-DIE[®], at the rate of 1 ounce per 50 square feet of surface area, into voids, cracks and crevices or other places where the insects are traveling or harboring. Apply a continuous visible film underneath and behind stairs, refrigerators and appliances. Treat on, around and behind moldings, shelving, baseboards, pipe openings, cracks, crevices, and other areas where insects hide. For severe infestations, apply TRI-DIE[®] behind cabinets and walls, in crawl spaces and attics at the rate of 16 ounces per 1000 square feet of surface area. Repeat treatment as necessary.

STORED PRODUCT PESTS: To aid in the control of Cadelles, Cheese Mites, Cigarette Beetles, Confused Flour Beetles, Dark Mealworms, Drugstore Beetles, Grain Mites, Granary Weevils, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles and Yellow Mealworms in mills, elevators and granaries. Clean up waste materials, dust, dirt, and any other debris. Thoroughly treat floors, walls and other surfaces in bins, storage and handling areas. Apply into cracks and crevices and other difficult to reach areas where insects may hide. This product may also be applied in voids or cracks and crevices in pantries and other infested areas within the structure.

FLEAS, LICE and TICKS: Treat pet bedding, floor and floor coverings and all areas where pests may inhabit. Treat cracks and crevices behind moldings, shelving, baseboards, pipe openings, and all other cracks. Repeat application when effectiveness diminishes.

BEDBUGS: Take bed apart. Dust into joints and channels. If hollow, such as square or round

tubing, see that the interior framework is well dusted. Mattresses may be dusted, especially tufts, folds, and edges. Picture frame moldings and all cracks and crevices in the room should be treated.

DRYWOOD TERMITES, POWDER POST BEETLES: To protect structures use an electric or hand operated rotary duster to distribute TRI-DIE[®] through insect entry points, such as, crawl holes into attics and crawl spaces of buildings, at a rate of 1 pound per 1000 square feet of floor area. All vents and blowers on the windward side should be closed so that all wood surfaces can be treated. In case of localized infestations, probe to locate galleries and drill holes to inject TRI-DIE[®] into the galleries and void areas. If a heavy infestation has developed, fumigation should be employed.

SOWBUGS, PILLBUGS, GROUND BEETLES, EARWIGS, MILLIPEDES, CENTIPEDES AND SCORPIONS: Apply in voids and cracks and crevices around doors and windows and other places where these pests may



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.



TRI-DIE® Silica & Pyrethrum Dust

enter premises. Treat behind baseboards, storage areas and other locations where these pests are found.

APPLICATIONS WITHIN FOOD AREAS

Food processing plants and warehouses, bakeries, beverage plants, restaurants, supermarkets, hotels, motels, hospitals and schools. Includes areas for receiving, storage (dry, cold, frozen or raw), packaging (canning, bottling, and grinding), enclosed processing systems (mills, dairies, and systems for processing edible oils and syrups).

LIMITED TO CRACK AND CREVICE®

TREATMENT ONLY: Using a bulbous duster apply a small amount of material directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housing, junction boxes or switch boxes, conduits or hollow equipment legs. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE® TREATMENT ARE NOT PERMITTED.

INDOOR AND OUTDOOR TREATMENTS

CARPENTER ANTS: Locate source of infestation or areas of suspected activity. Treat insect tunnels and cavities and void areas. Treat points of entry or exits from voids such as around doors, windows and siding.

BOXELDER BUGS, CLOVER MITES, CLUSTER FLIES AND ELMLEAF BEETLES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames. Repeat as necessary.

BEEES AND WASPS (in Ground or Voids): If practical it is advisable to treat wasp and bee nests in the evening when insects are

less active and have returned to the nest. Using hand or power duster or other suitable means, with extension tubes if necessary, thoroughly dust nest and entrance and surrounding areas where insects alight. Locate nests in wall voids. If necessary, drill a hole in the area. Apply dust in void and reseal. For best results check nests carefully one or two days after treatment to ensure complete kill. Repeat if necessary. Remove or destroy nest to prevent emergence of newly-hatched insects.

SEWER TREATMENTS: Cockroach or other insect infestations in sanitary sewers may be treated with approximately 6 ounces of TRI-DIE® per manhole. When treating sewers, one manhole per 200 feet of line should be treated. Apply product to sides of manhole and other surfaces with hand or power duster. Treatments may also be made by removing the manhole cover and replacing with a disc of plywood bored to receive the usual flexible hose from portable or truck-mounted ventilating equipment, 6 ounces of TRI-DIE® is introduced into the ventilating air to be dispersed quickly and thoroughly through the system.

OUTDOOR TREATMENTS

ANTS, CENTIPEDES, CLOVER MITES, COCKROACHES, CRICKETS, EARWIGS, GROUND BEETLES, MILLIPEDES, PILLBUGS, SCORPIONS, SOWBUGS AND SPIDERS: Treat cracks around doors, windows, under siding and voids foundation level, beneath guards on loading docks, into cracks in foundations and floors and in other places where these pests may enter building. Repeat application as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Keep away from moisture. Sweep up and dispose of with chemical waste.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or an approval waste disposal facility.

CONTAINER DISPOSAL: Completely empty liner by shaking and taping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting unless directed by a physician. Contains petroleum solvent. Contact a physician or Poison Control Center immediately.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation/symptoms persist.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Special precautions should be taken so that all residue will be out of reach of children or domestic animals. Avoid contamination of food and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate.

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682

© 2000 Whitmire Micro-Gen Research Laboratories, Inc.

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.

Revised: 11/00